

Contents

| | |
|---|-------------|
| Acknowledgements | ii |
| Preface | v |
| Acronyms | vi |
| Technical summary | xvii |
| 1. Introduction | 1 |
| 1.1 Background | 1 |
| 1.2 Large marine ecosystems | 2 |
| 1.3 The Western Pacific Warm Pool | 3 |
| References | 5 |
| 2. Large marine ecosystems assessment methodology | 11 |
| 2.1 Approach to assessment and management of LMEs: LME modules | 11 |
| 2.2 Conceptual framework | 11 |
| 2.3 Scale and scope of the assessment | 14 |
| 2.4 Approach to the comparative assessment | 14 |
| 2.5 Assessment process | 15 |
| 2.6 Organization of this report | 15 |
| References | 16 |
| 3. Socio-economics: Examining socio-economic dimensions of risk and vulnerability among coastal inhabitants of large marine ecosystems | 21 |
| 3.1 Introduction | 22 |
| 3.2 Findings | 23 |
| 3.3 Discussion and conclusions | 41 |
| 3.4 Methodology and analysis | 43 |
| References | 53 |
| 4. Governance: Assessment of governance arrangements for transboundary large marine ecosystems ... | 65 |
| 4.1 Introduction | 66 |
| 4.2 Main findings and discussion | 69 |
| 4.3 Conclusion | 82 |
| 4.4 Methodology and analysis | 83 |
| References | 85 |
| 5. Productivity | 91 |
| 5.1 Primary productivity patterns and trends | 91 |
| 5.1.1 Introduction | 92 |
| 5.1.2 Data and methodologies | 92 |
| 5.1.3 Major findings, discussion, and conclusions | 93 |
| 5.2 Sea surface temperature trends in large marine ecosystems | 101 |
| 5.2.1 Introduction | 101 |
| 5.2.2 Main findings, discussion and conclusions | 102 |
| 5.2.3 Data and methodology | 108 |
| References | 109 |

| | | |
|-----------|---|------------|
| 6. | Fish and Fisheries | 113 |
| 6.1 | The Status of fisheries in large marine ecosystems, 1950–2010 | 113 |
| 6.1.1 | Introduction | 115 |
| 6.1.2 | Main findings | 116 |
| 6.1.3 | Discussion | 125 |
| 6.1.4 | Methodology and analysis | 126 |
| 6.1.5 | The Western Pacific Warm Pool | 130 |
| | References | 134 |
| 6.2 | Fishery production potential of large marine ecosystems: a prototype analysis | 139 |
| 6.2.1 | Introduction | 139 |
| 6.2.2 | Methods and data sources | 140 |
| 6.2.3 | Results | 143 |
| 6.2.4 | Discussion | 146 |
| | References | 149 |
| 7. | Pollution and Ecosystem Health | 153 |
| 7.1 | Floating plastic debris | 153 |
| 7.1.1 | Introduction | 154 |
| 7.1.2 | Main findings, discussion, and conclusions | 155 |
| 7.1.3 | Methodology and analysis | 162 |
| | References | 163 |
| 7.2 | Pollution status of persistent organic pollutants | 165 |
| 7.2.1 | Introduction | 166 |
| 7.2.2 | Findings, discussion, and conclusions | 167 |
| 7.2.3 | Methodology and analysis | 173 |
| | References | 176 |
| 7.3 | Nutrient inputs from river systems to coastal waters | 179 |
| 7.3.1 | Introduction | 179 |
| 7.3.2 | Findings, discussion, and conclusions | 180 |
| 7.3.3 | Methodology and analysis | 187 |
| | References | 192 |
| 7.4 | Extent of mangroves and drivers of change | 195 |
| 7.4.1 | Introduction | 195 |
| 7.4.2 | Findings and discussion | 197 |
| 7.4.3 | Methodology | 198 |
| 7.4.4 | Limitations | 200 |
| | References | 201 |
| 7.5 | Reefs at Risk Index | 203 |
| 7.5.1 | Introduction | 204 |
| 7.5.2 | Findings and discussion | 205 |
| 7.5.3 | Methodology | 210 |
| | References | 213 |

| | | |
|-----------|---|------------|
| 7.6 | Change in protected area coverage within large marine ecosystems | 215 |
| 7.6.1 | Introduction | 215 |
| 7.6.2 | Findings and discussion | 216 |
| 7.6.3 | Methodology | 218 |
| | References | 221 |
| 7.7 | Cumulative human impacts in the world's large marine ecosystems | 223 |
| 7.7.1 | Introduction | 224 |
| 7.7.2 | Findings, discussion, and conclusions | 225 |
| 7.7.3 | Methodology and analysis | 229 |
| | References | 232 |
| 7.8 | Ocean Health Index for the worlds large marine ecosystems | 237 |
| 7.8.1 | Introduction | 238 |
| 7.8.2 | Findings, discussion, and conclusions | 239 |
| 7.8.3 | Methodology and analysis | 242 |
| | References | 244 |
| 8. | Identifying patterns of risk among large marine ecosystems using multiple indicators | 251 |
| 8.1 | Introduction | 253 |
| 8.2 | Results, discussion, and conclusions | 254 |
| 8.3 | Methodology and analysis | 267 |
| | References | 271 |
| 9. | Conclusion | 287 |
| 9.1 | Meeting Transboundary Waters Assessment Programme (TWAP) objectives | 287 |
| 9.2 | Main conclusions | 287 |
| 9.3 | Patterns of risk among LMEs based on multiple indicators | 291 |
| 9.4 | Overall conclusion | 291 |
| 9.5 | Target audience | 292 |
| 9.6 | Future TWAP LME assessments | 292 |
| | Glossary | 294 |
| | References | 299 |

Figures

| | | |
|-------------|---|-----|
| Figure 1.1 | Map of the 66 Large marine ecosystems of the world, plus location of the Western Pacific Warm Pool | 2 |
| Figure 1.2 | Warm-water areas of the global ocean, highlighting the Western Pacific Warm Pool | 4 |
| Figure 2.1 | LME assessment conceptual framework illustrating the interactions between the human and natural systems | 13 |
| Figure 3.1 | The size of coastal populations – a proxy measure of pressure on ecosystem services provided by LMEs | 24 |
| Figure 3.2. | Proportion of coastal zone population that is rural | 25 |
| Figure 3.3 | Populations below national poverty lines – an indicator of socio-economic vulnerability. | 26 |
| Figure 3.4 | Coastal eutrophication potential and the Human Development Index | 28 |
| Figure 3.5 | Night light distribution as a spatial proxy for level of economic development (Night Light Development Index) | 29 |
| Figure 3.6 | Night Light Development Index paired with the Human Development Index | 29 |
| Figure 3.7 | Human Development Index for LMEs | 30 |
| Figure 3.8 | Shared Socio-economic Pathways | 31 |
| Figure 3.9 | The use of Representative Concentration Pathways and Shared Socio-economic Pathways in the analysis of impact, adaptation, and vulnerability in relation to global environmental change | 32 |
| Figure 3.10 | Human Development Index in 2100 based on two Shared Socio-economic Pathways. | 33 |
| Figure 3.11 | Present-day Climate-related Extreme Events Threat Index | 34 |
| Figure 3.12 | Projected mean sea surface height in 2100 and Sea-level Rise Threat Index in 2100 for two shared Socio-economic Pathways | 36 |
| Figure 3.13 | Contemporary Threat Index | 39 |
| Figure 4.1 | Frequency distribution of LMEs by number of transboundary issues (including the WPWP) | 71 |
| Figure 4.2 | Frequency distribution of LMEs by average per cent completeness of all arrangements in each LME (including the WPWP). | 72 |
| Figure 4.3 | Global distribution of levels of completeness and perceived risk for 49 transboundary LMEs and the WPWP | 73 |
| Figure 4.4 | Completeness distribution of fisheries, pollution, and biodiversity arrangements across all LMEs and the WPWP | 74 |
| Figure 4.5 | The distribution of scores for each of the seven policy-cycle stages for arrangements addressing fisheries, pollution, and biodiversity across all LMEs and the WPWP | 75 |
| Figure 4.6 | Distribution of integration scores for LMEs (including the WPWP) | 76 |
| Figure 4.7 | Global distribution of levels of integration and perceived risk for 49 transboundary LMEs and the WPWP | 77 |
| Figure 4.8 | Level of overall country engagement in binding and non-binding agreements by number of LMEs (including the WPWP). | 79 |
| Figure 4.9 | Global distribution of levels of engagement and perceived risk for 49 transboundary LMEs and the WPWP | 80 |
| Figure 4.10 | Per cent engagement by countries in binding and non-binding agreements for transboundary issues | 80 |
| Figure 5.1 | Distribution of average surface chlorophyll <i>a</i> throughout the global ocean, 1998–2013. | 94 |
| Figure 5.2 | Distribution of average annual primary productivity throughout the global ocean, 1998–2013. | 95 |
| Figure 5.3 | Classification of 66 LMEs and the WPWP into five groups by productivity | 95 |
| Figure 5.4 | Global distribution of the five productivity classification groups for 66 LMEs and the WPWP | 96 |
| Figure 5.5 | Trends in chlorophyll <i>a</i> (2003–2013) in relation to latitude | 97 |
| Figure 5.6 | Global distribution of chlorophyll <i>a</i> samples, 1998–2013. | 97 |
| Figure 5.7 | Relationship of LME sampling frequency and coverage, 1998–2013 | 98 |
| Figure 5.8 | Long-term sea surface temperature trends (net changes) in 66 LMEs, 1957–2012 | 104 |
| Figure 5.9 | Sea surface temperature time series in selected LMEs of the North Pacific | 105 |
| Figure 5.10 | Sea surface temperature time series in selected LMEs of the Western North Atlantic | 106 |

| | | |
|-------------|--|-----|
| Figure 5.11 | Sea surface temperature time series in selected LMEs of the Eastern North Atlantic (European seas) | 107 |
| Figure 6.1 | Time series of landings by species in all LMEs, 1950–2010 | 116 |
| Figure 6.2 | Percentage of the total annual catch from bottom-impacting gears in all LMEs, 1950–2010 | 117 |
| Figure 6.3 | Ex-vessel value of reported landings in all LMEs, by country | 118 |
| Figure 6.4 | Aggregate effective fishing effort in LMEs, 1950–2006 | 119 |
| Figure 6.5 | Paired stock status plots for the catch of all LMEs, assessing the status of stocks defined as taxa with a time series of landings in an LME | 120 |
| Figure 6.6 | Primary production required (PPR) to sustain fisheries in the world's LMEs, an expression of their 'ecological footprint' | 122 |
| Figure 6.7 | Two indicators based on the trophic levels of exploited fish, used to characterize the fisheries in the LMEs | 123 |
| Figure 6.8 | The Western Pacific Warm Pool and the EEZs of the countries that it includes | 130 |
| Figure 6.9 | Reported landings in the WPWP from 1950 to 2010, based on FAO data spatially allocated | 131 |
| Figure 6.10 | Fisheries-related indicators for the WPWP, 1950–2010 | 132 |
| Figure 6.11 | Food-web structure employed in this analysis | 141 |
| Figure 6.12 | Distribution patterns for total chlorophyll <i>a</i> and primary production | 143 |
| Figure 6.13 | Estimated production levels in the absence of exploitation, by functional group, for LMEs represented in this study | 144 |
| Figure 6.14 | Ratio of landings to phytoplankton primary production | 145 |
| Figure 7.1 | World plastics production, 1950–2012, showing the rapid increase in production to match demand | 154 |
| Figure 7.2 | Relative abundance of floating micro-plastics in 66 LMEs, based on model estimates and placed into five risk categories | 156 |
| Figure 7.3 | Relative abundance of floating macro-plastics in 66 LMEs, based on model estimates, and placed into five risk categories | 157 |
| Figure 7.4 | Spatial distribution of the relative abundance of floating (a) micro-plastics and (b) macro-plastics in 66 LMEs, based on model estimates | 159 |
| Figure 7.5 | Floating macro-plastics in the Straits of Malacca (eastern extremity of Bay of Bengal LME, 'highest' risk category) | 160 |
| Figure 7.6 | Floating micro- and macro-plastics in the North Sea LME ('high' risk category), retrieved from the stomach of an open-water foraging bird, the northern fulmar (<i>Fulmarus glacialis</i>) | 161 |
| Figure 7.7 | Correlation between median PCB concentrations in plastic resin pellets and in mussels | 168 |
| Figure 7.8 | Concentrations of PCBs in beached plastic pellets within LMEs | 169 |
| Figure 7.9 | Concentrations of DDTs in beached plastic pellets within LMEs | 169 |
| Figure 7.10 | Concentrations of HCHs in beached plastic pellets within LMEs | 170 |
| Figure 7.11 | Nitrogen load risk categories for LMEs for a) 2000, b) 2030, and c) 2050 | 181 |
| Figure 7.12 | Index of Coastal Eutrophication Potential (ICEP) risk categories for LMEs for a) 2000, b) 2030, and c) 2050 | 182 |
| Figure 7.13 | Merged nutrient risk categories for LMEs for a) 2000, b) 2030, and c) 2050 | 184 |
| Figure 7.14 | Dissolved inorganic nitrogen (DIN) yield, Index of Coastal Eutrophication Potential (ICEP), and sources of DIN in river basins draining to the Bay of Bengal LME | 186 |
| Figure 7.15 | Conceptual diagram of the Global NEWS model construction, sub-models, and parameters | 190 |
| Figure 7.16 | Mangrove areas within LMEs | 197 |
| Figure 7.17 | Mangrove extent within each LME, and the WPWP, expressed as a) per cent cover and b) area | 198 |
| Figure 7.18 | LME regions extended to incorporate inland mangrove areas into the analysis | 200 |
| Figure 7.19 | Integrated Local Threat Index scores for LMEs containing coral reefs and the WPWP, shown by risk category | 206 |
| Figure 7.20 | Proportion of LME and WPWP coral reef extent by threat level for a) present-day integrated local threats and b) present-day integrated local threats and past thermal stress | 207 |
| Figure 7.21 | Projected proportion of LMEs and WPWP coral reef area by threat level for global threats (warming and acidification) by a) 2030 and b) 2050 | 208 |
| Figure 7.22 | Warm-water coral reef areas within LMEs and the WPWP | 209 |

| | |
|---|-----|
| Figure 7.23 Coral cover within each LME and the WPWP shown as a) percentage and b) area | 209 |
| Figure 7.24 LMEs extended inland to incorporate all coral reef areas within LMEs into the analysis | 210 |
| Figure 7.25 Marine Protected Areas designated by a) 1982 and b) 2014 | 216 |
| Figure 7.26 Index of percentage change (1982–2014) in total area covered by MPAs per LME | 217 |
| Figure 7.27 Cumulative area of MPAs in all LMEs and the WPWP | 218 |
| Figure 7.28 Cumulative human impact risk categories of LMEs and the WPWP. | 225 |
| Figure 7.29 Ocean Health Index score by LME. | 239 |
| Figure 7.30 Ocean Health Index risk category by LME. | 240 |
| Figure 8.1 LME clusters and their distinguishing features. | 256 |
| Figure 8.2 Heat map of indicator scores, risk categories of governance metrics, and the Ocean Health Index. | 257 |
| Figure 8.3 Maps of the three main principal components showing risks related to combinations of indicators. | 260 |
| Figure 8.4 Risk assessments based on averages of the standardized indicator values for each module. | 263 |
| Figure 8.5 TWAP risk scores by LME, averaged across modules, compared with OHI risk scores | 264 |

Annex Figures

| | |
|---|-----|
| Annex Figure 8-A Dendrogram illustrating the six clusters. | 272 |
| Annex Figure 8-B PCA biplot illustrating LME clusters in relation to the 11 indicator axes. | 273 |

Tables

| | | |
|------------|--|-----|
| Table 3.1 | Comparison of Present-day Climate-related Extreme Events Threat Index values with and without the HDI Gap as a vulnerability measure for the two highest risk categories | 34 |
| Table 3.2 | Contemporary Threat Index for each LME | 38 |
| Table 3.3 | Average values of attributes of LMEs by risk category of Contemporary Threat Index. | 39 |
| Table 3.4 | Assessment of Pacific island states within the Western Pacific Warm Pool. | 40 |
| Table 4.1 | Risk categories and ranges for the three indicators. | 69 |
| Table 4.2 | Number of arrangements, and scores and risk levels for completeness, integration, and engagement indicators for transboundary LMEs and the WPWP. | 69 |
| Table 4.3 | Steps required to assess governance architecture in a system to be governed | 83 |
| Table 4.4 | Scoring criteria for policy-cycle stages for each arrangement. | 83 |
| Table 5.1 | Comparison between <i>in situ</i> and satellite-based estimates of primary productivity for the Northeast US Continental Shelf LME. | 93 |
| Table 5.2 | Net sea surface temperature changes in LMEs and the WPWP, 1957–2012. | 103 |
| Table 5.3 | Classification of LMEs based on net change in sea surface temperature, 1957–2012 | 106 |
| Table 6.1 | Five relative risk categories and cut-off points used for grouping the LMEs for each indicator. | 118 |
| Table 6.2 | Fisheries-based indicators of the WPWP compared to those of the LMEs | 133 |
| Table 6.3 | Definitions of functional groups for this analysis | 141 |
| Table 7.1 | Main polymer types, typical applications, and specific gravity | 155 |
| Table 7.2 | Grouping of LMEs into five categories on the basis of the relative concentration of (a) micro-plastics and (b) macro-plastics, based on model estimates | 158 |
| Table 7.3 | Risk categories for concentrations of (a) PCBs, (b) DDTs, and (c) HCHs in plastic pellets in LMEs | 175 |
| Table 7.4 | LMEs with risk category changes between 2000 and 2030, and between 2030 and 2050. | 183 |
| Table 7.5 | Risk categories for the nitrogen load sub-indicator | 188 |
| Table 7.6 | Risk categories for the Index of Coastal Eutrophication Potential (ICEP) sub-indicator | 189 |
| Table 7.7 | Risk categories for the merged nutrient risk indicator. | 189 |
| Table 7.8 | Input data sets used in the Global NEWS model for nitrogen and phosphorus | 190 |
| Table 7.9 | Relative importance of drivers of mangrove loss and their projected changes, by region. | 199 |
| Table 7.10 | Risk categories based on Integrated Local Threat Index scores for LMEs containing coral reefs and the WPWP. | 205 |
| Table 7.11 | Example of how the total integrated threat scores were calculated for three of the LMEs. | 211 |
| Table 7.12 | Threat level categories based on change in MPA coverage in LMEs. | 221 |
| Table 7.13 | Summary of data layers used to calculate CHI | 230 |
| Table 7.14 | Summary of CHI values per risk category for global LMEs. | 231 |
| Table 7.15 | Definitions of the goals and sub-goals of the Ocean Health Index. | 240 |
| Table 7.16 | Summary of Ocean Health Index values per risk category for all LMEs | 243 |
| Table 8.1 | Indicators selected for the multivariate analysis | 254 |
| Table 8.2 | PCA loadings on each of the eleven principal components and proportion of variance explained by each component | 259 |

Annex Tables

| | | |
|-----------------|--|-----|
| Annex Table 1-A | LMEs and Western Pacific Warm Pool and bordering countries | 6 |
| Annex Table 2-A | TWAP LMEs assessment working group of institutional partners and experts | 17 |
| Annex Table 3-A | Socio-economic indicators, methods, and data sources used in assessing risk and vulnerability of coastal populations next to LMEs | 54 |
| Annex Table 3-B | Fishing and tourism revenues | 57 |
| Annex Table 3-C | A comparison of the elements of Shared Socio-economic Pathways 1 and 3 used in the study | 59 |
| Annex Table 3-D | Comparing coastal populations projected by the UNDP and population estimates based on Shared Socio-economic Pathways | 63 |
| Annex Table 4-A | Breakdown of transboundary issues for transboundary LMEs and the WPWP | 90 |
| Annex Table 6-A | Classifying the 64 assessed LMEs into 5 relative risk categories for each fisheries-related indicator | 136 |
| Annex Table 7-A | Full results of CHI and individual stressor impact scores for each LME and the WPWP | 233 |
| Annex Table 7-B | Full results of Ocean Health Index scores and component goal scores for each LME | 245 |
| Annex Table 8-A | Forty-eight indicators used in assessing LMEs | 274 |
| Annex Table 8-B | Correlation matrix | 275 |
| Annex Table 8-C | Variance matrix | 276 |
| Annex Table 8-D | Average scores of selected indicators, by module, calculation of the TWAP risk score, and comparison of risk ranks based on the LME TWAP risk score and the Ocean Health Index | 277 |
| Annex Table 8-E | TWAP risk score calculations and risk rank comparisons for the 47 inhabited transboundary LMEs | 279 |
| Annex Table 8-F | Raw unstandardized values for the 11 chosen indicators | 281 |
| Annex Table 8-G | Standardized and rescaled values for the 11 chosen indicators | 283 |

Boxes

| | | |
|---------|---|-----|
| Box 4.1 | Explanation of key governance terms used in this chapter | 67 |
| Box 6.1 | Sea Around Us fisheries indicators: definitions and interpretation | 115 |
| Box 7.1 | An example of within-LME variation of nutrient loads and sources: the Bay of Bengal LME | 186 |
| Box 8.1 | Overview of statistical techniques | 255 |